

AFFINE TRANSFORMATION WITH ZERO ENTROPY AND NILSYSTEMS

ZHENGXING LIAN

ABSTRACT. In this paper, we study affine transformations on tori, nilmanifolds and compact abelian groups. For these systems, we show that an equivalent condition for zero entropy is the orbit closure of each point has a nice structure. To be precise, the affine systems on those spaces are zero entropy if and only if the orbit closure of each point is isomorphic to an inverse limit of nilsystems.

1. INTRODUCTION

By a $(\mathbb{Z}$ -action) topological dynamical system, one means a pair (X, T) , where X is a compact metric space and T is a continuous selfmap of X . In this paper, we consider the cases that X is a torus, a nilmanifold or a compact abelian group.

For a topological group G and a cocompact subgroup Γ , an *affine transformation* τ on G/Γ is defined by an Γ -invariant automorphism A of G and a fixed element $g_0 \in G$, i.e. $\tau(g\Gamma) = g_0 A(g)\Gamma$. In particular, when G itself is a compact group and Γ is trivial, the affine transformation $\tau : G \rightarrow G$ is the map $\tau(g) = g_0 A(g)$, where $g_0 \in G, A \in \text{Aut}(G)$.

In this paper, we study affine transformations on tori, nilmanifolds and compact abelian groups, and give an equivalent condition of zero entropy for these systems. Our main result is as follows:

Theorem 1.1. *Let (X, τ) be one affine transformation of the following spaces:*

- (1) *tori;*
- (2) *nilpotent manifolds;*
- (3) *compact abelian groups.*

Then (X, τ) is zero entropy if and only if for each $x \in X$, the closure of the orbit of x with τ $(\overline{\text{orb}(x, \tau)}, \tau)$ is isomorphic to an inverse limit of nilsystems. We might call the system $(\overline{\text{orb}(x, \tau)}, \tau)$ the orbit system.

As an application, we show Sarnak conjecture holds for the zero entropy affine transformations on tori, nilmanifolds and compact abelian groups, which is first proved in [17]. According to Green and Tao's result[8](i.e. *Möbius function* is orthogonal with nilsequence), we give different methods to show this.

The paper is organized as follows: In Section 2, we introduce some basic conceptions and results needed in this paper. In Section 3, we prove Theorem 1.1 by three

2010 *Mathematics Subject Classification.* Primary: 37B05.

Key words and phrases. Affine transformations; nilsequences; entropy.

The author is supported by NNSF of China (11171320, 11571335).

different cases. To prove this theorem, we try to analysis the returning time of each point of X . Then we use some theorems in [16] to prove our results.

Acknowledgments. The author would like to thank Wen Huang, Song Shao and Xiangdong Ye for introducing him this topic and their constant support.

2. PRELIMINARIES

In this section, we introduce some basic definitions and results needed in this paper.

2.1. Topology entropy. Let (X, τ) be a topological dynamical system, where $\tau : X \rightarrow X$ is continuous, X is a compact metric space with metric ρ . Let $\rho_n(x, y) = \max_{0 \leq i \leq n-1} \rho(\tau^i x, \tau^i y)$. A set $E \subset X$ is (n, ϵ) -separated if for any $x \neq y \in E$, $\rho_n(x, y) > \epsilon$. Denote by $N(n, \epsilon)$ the maximum cardinality of an (n, ϵ) -separated set. The *topological entropy* of (X, τ) is defined by

$$h_{top}(X, \tau) = \lim_{\epsilon \rightarrow 0} (\limsup_{n \rightarrow \infty} \frac{1}{n} \log N(n, \epsilon))$$

Let $\pi : (X, \tau) \rightarrow (Y, S)$ be a factor map, i.e. π is a continuous surjection and $\pi \circ \tau = S \circ \pi$. Then $h_{top}(X, \tau) \geq h_{top}(Y, S)$.

In this paper, we will need the following famous result:

Theorem 2.1. [19] *Let D be a homomorphism of a d -dim torus. $\{\lambda_1, \dots, \lambda_d\}$ are the eigenvalues of D . Then the entropy of D is:*

$$h(\mathbb{T}^d, D) = \sum_{j=1}^d \log |\lambda_j|_+,$$

where $\log |\lambda_j|_+ = \max\{\log |\lambda_j|, 0\}$.

2.2. Nilsystem.

2.2.1. Nilpotent groups. Let G be a group. For $g, h \in G$, we write $[g, h] = ghg^{-1}h^{-1}$ for the commutator of g and h and we write $[A, B]$ for the subgroup spanned by $\{[a, b] : a \in A, b \in B\}$. The commutator subgroup $G_j, j \geq 1$, are defined inductively by setting $G_1 = G$ and $G_{j+1} = [G_j, G]$. Let $d \geq 1$ be an integer. We say that G is *d-step nilpotent* if G_{d+1} is the trivial subgroup.

2.2.2. Nilmanifolds. Let G be a d -step nilpotent Lie group and Γ is a discrete cocompact subgroup of G , i.e. a uniform subgroup of G . The compact manifold $X = G/\Gamma$ is called a *d-step nilmanifold*. The group G acts on X by left translations and we write this action as $(g, x) \rightarrow gx$. The Haar measure μ of X is the unique probability measure on X invariant under this action. Let $g \in G$ and τ be the transformation $\tau(x) \rightarrow gx$ of X . Then (X, τ, μ) is called a *d-step nilsystem*. See [4] for more details.

2.2.3. Systems of order d . We also make use of inverse limits of nilsystems and so we recall the definition of an inverse limit of systems (restricting ourselves to the case of sequential inverse limits). If $(X_i, \tau_i)_{i \in \mathbb{N}}$ are systems with $\text{diam}(X_i) \leq M \leq \infty$ and $\phi_i : X_{i+1} \rightarrow X_i$ are factor maps, the *inverse limit* of the systems is defined to be the compact subset of $\prod_{i \in \mathbb{N}} X_i$ given by $\{(x_i)_{i \in \mathbb{N}} : \phi_i(x_{i+1}) = x_i, i \in \mathbb{N}\}$, which is denoted by $\lim_{\leftarrow} \{X_i\}_{i \in \mathbb{N}}$. It is a compact metric space endowed with the distance $\rho(x, y) = \sum_{i \in \mathbb{N}} \frac{1}{2^i} \rho_i(x_i, y_i)$, where ρ_i is the distance of (X_i, τ_i) . We note that the maps $\{\tau_i\}$ induce a transformation τ on the inverse limit.

Definition 2.2. [14] A system (X, τ) is called a *system of order d* , if it is an inverse limit of d -step minimal nilsystems.

2.3. Regionally proximal relation. Let (X, τ) be a topological dynamical system. The *regionally proximal relation* $RP = RP(X, \tau) \subset X \times X$ is the set which contains all the point (x, y) if there are sequences $x_i, y_i \in X, n_i \in \mathbb{Z}$ such that $x_i \rightarrow x, y_i \rightarrow y$ and $(\tau \times \tau)^{n_i}(x_i, y_i) \rightarrow (z, z), i \rightarrow \infty$, for some $z \in X$.

Definition 2.3. [13] Let (X, τ) be a topological dynamical system with metric ρ . Let $d \geq 1$ be an integer. A pair $(x, y) \in X \times X$ is said to be *regionally proximal of order d* if for any $\delta > 0$, there exist $x', y' \in X$ and a vector $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$ such that $\rho(x, x') < \delta, \rho(y, y') < \delta$, and

$$\rho(\tau^{n \cdot \epsilon} x', \tau^{n \cdot \epsilon} y') < \delta \text{ for any } \epsilon \in \{0, 1\}^d \setminus \{(1, 1, \dots, 1)\},$$

where $n \cdot \epsilon = n_1 \epsilon_1 + \dots + n_d \epsilon_d$. The set of regionally proximal pairs of order d is denoted by $RP^{[d]}(X)$.

It is easy to see that $RP^{[d]}$ is a closed and invariant equivalence relation.

Theorem 2.4. [14, 16] *Let (X, τ) be a minimal topological dynamical system and $d \in \mathbb{N}$. Then $(X/RP^{[d]}, \tau)$ is the maximal d -step nilfactor of (X, τ) .*

It means that $(X/RP^{[d]}, \tau)$ is a system of order d . Also, any system of order d , which is a factor of (X, τ) , is a factor of $(X/RP^{[d]}, \tau)$.

Now we use $RP^{[d]}$ to define ∞ -step nilsystem[5]. Observe that

$$\dots \subset RP^{[d+1]} \subset RP^{[d]} \subset \dots \subset RP^{[1]} = RP(X, \tau)$$

Let $RP^{[\infty]} = \bigcap_{d \geq 1} RP^{[d]}$. It follows that for any minimal system (X, τ) , $RP^{[\infty]}$ is a closed invariant equivalence relation.

Definition 2.5. A minimal system (X, τ) is an *∞ -step nilsystem* or a *system of order ∞* , if the equivalence relation $RP^{[\infty]}$ is trivial. i.e., coincides with the diagonal.

2.4. Almost automorphic. In [16], the authors gave some relationship between nilsystems and almost automorphic points.

For a topological dynamical system (X, τ) , a point $x \in X$ is said to be *almost automorphic* (AA for short) if from any sequence $\{n'_i\} \subset \mathbb{Z}$ one may extract a subsequence $\{n_i\}$ such that

$$\lim_{j \rightarrow \infty} \lim_{i \rightarrow \infty} \tau^{n_i - n_j} x = x.$$

A point $x \in X$ is called *d -step almost automorphic* (or *d -step AA* for short) if

$$RP^{[d]}[x] = \{y \in X : (x, y) \in RP^{[d]}\} = \{x\}.$$

A minimal topological dynamical system is called *d-step almost automorphic* if it has a *d*-step AA point. Since $RP^{[d]}$ is an equivalence relation for minimal topological dynamical system, by definition it follows that for a minimal system (X, τ) , it is a *d*-step AA system for some $d \in \mathbb{N}$ if and only if it is an almost one-to-one extension of its maximal *d*-step nilfactor.

In [16], one also describes AA point by the returning time of one point to its open neighborhoods.

2.5. Nil Bohr sets. A subset A of $\mathbb{Z}$ is a *Bohr-set* if there exist $m \in \mathbb{N}$, $\alpha \in \mathbb{T}^m$ and a non-empty open set $U \subset \mathbb{T}^m$ such that $\{n \in \mathbb{Z} : n\alpha \in U\}$ is contained in A . The set A is a *Bohr₀-set* if additionally $0 \in U$.

A subset A of $\mathbb{Z}$ is a *Nil_d Bohr₀ set* if there exist a *d*-step nilsystem (X, T) , $x_0 \in X$ and an open neighborhood U of x_0 such that $N(x_0, U) =: \{n \in \mathbb{Z} : T^n x_0 \in U\}$ is contained in A . Denote by $\mathcal{F}_{d,0}$ the family consisting of all *Nil_d Bohr₀-sets* [12].

Let $\mathcal{F}_{GP_d}$ be the family generated by the set of forms

$$\bigcap_{i=1}^k \{n \in \mathbb{Z} : P_i(n) \pmod{\mathbb{Z}} \in (-\epsilon_i, \epsilon_i)\}$$

where $k \in \mathbb{N}$, $P_1, \dots, P_k$ are generalized polynomials of degree $\leq d$, and $\epsilon_i > 0$.

In this paper, we consider generalized polynomial family $GP = \bigcup_{d=0}^{\infty} GP_d$, which is defined by a sequence of family GP_d . Where GP_d is the smallest family satisfies:

- (1) Any k , let $p_i \in \mathbb{N}$, $0 \leq i \leq k$. If $f_1, f_2, \dots, f_l$ with $f_i \in GP_{p_i}$ and $\sum_{i=0}^l p_i \leq d$, then $an^{p_0} \lceil f_1 \rceil \lceil f_2 \rceil \dots \lceil f_k \rceil \in GP_d$. Where $\lceil x \rceil$ is the smallest integer such that $\lceil x \rceil \geq x$.
- (2) If $g, h \in GP_d$, then $gh, g \pm h, cg, \lceil g \rceil \in GP_d$. Where $c \in \mathbb{R}$

And $G_0 = \mathbb{R}$.

If $f \in GP_d$, we say the degree of f $\deg(f) \leq d$.

Theorem 2.6. [16] *Let $d \in \mathbb{N}$. Then $\mathcal{F}_{d,0} = \mathcal{F}_{GP_d}$.*

Theorem 2.7. [16] *Let (X, T) be a minimal topological dynamical system, $x \in X$ and $d \in \mathbb{N} \cup \{\infty\}$. Then the following statements are equivalent:*

- (1) x is *d*-step AA point.
- (2) $N(x, V) \in \mathcal{F}_{d,0}$ for each neighborhood V of x .

3. THE PROOF OF THEOREM 1.1

Let (X, τ) be one affine transformation of the following spaces:

- (1) tori;
- (2) nilpotent manifolds;
- (3) compact abelian groups.

We want to show that (X, τ) is zero entropy if and only if for each $x \in X$, the orbit system $(\overline{\text{orb}(x, \tau)}, \tau)$ is isomorphic to an inverse limit of nilsystems. The difficult part is to show if (X, τ) is zero entropy, then for each $x \in X$, the orbit system $(\overline{\text{orb}(x, \tau)}, \tau)$ is isomorphic to an inverse limit of nilsystems. Since the proofs for tori, nilpotent manifolds, and compact abelian groups are quite different, we will

prove them separately. Before that, we show the easy direction of the proof of Theorem 1.1.

3.1. The easy part of the proof of Theorem 1.1.

Now we assume that for each $x \in X$, the orbit system $(\overline{\text{orb}(x, \tau)}, \tau)$ is isomorphic to an inverse limit of nilsystems. We will show that the entropy of the system is zero. In fact, we show that (X, τ) is distal. Since each distal system has zero entropy, we finish the proof.

In fact, if for each point $x \in X$, the closure of $\text{orb}(x, \tau)$ is an inverse limit of nilsystems, then X should be distal. That is, if $y, y' \in \overline{\text{orb}(x, \tau)}$, since $(\overline{\text{orb}(x, \tau)}, \tau)$ is distal, then (y, y') is distal.

If $y \in \overline{\text{orb}(x, \tau)}$ and $y' \in \overline{\text{orb}(x', \tau)}$ for two different orbits with $\overline{\text{orb}(x, \tau)} \cap \overline{\text{orb}(x', \tau)} = \emptyset$, since $\overline{\text{orb}(x, \tau)}$ and $\overline{\text{orb}(x', \tau)}$ is closed, $\rho(y, y') \geq \min\{\rho(z, z') \mid z \in \overline{\text{orb}(x, \tau)}, z' \in \overline{\text{orb}(x', \tau)}\} = c_{x, x'} > 0$ for some fix number $c_{x, x'}$. Then (X, τ) is distal.

3.2. Results on tori and nilpotent manifolds. The main aim of this section is to prove the following results:

Theorem 3.1. *Let (X, τ) be an affine transformation with zero entropy. If X is one of the following spaces:*

- (1) *tori;*
- (2) *nilpotent manifolds;*

then there is a fixed d , such that for each $x \in X$, an open neighborhood U of x and its return time $N(g, U) = \{n : \tau^n x \in U\}$, we have $N(x, U) \supset \bigsqcup_{j=1}^b N_j$ ($\bigsqcup$ means disjoint union) for some $b \in \mathbb{N}$ with $N_j \in \mathcal{F}_{GP_d}$ and $N_i \cap N_j = \emptyset$ for each $i \neq j$.

Theorem 3.2. *Let (X, τ) be an affine transformation with zero entropy. If X is one of the following spaces:*

- (1) *tori;*
- (2) *nilpotent manifolds;*

then for each $g \in G$, the orbit system $(\overline{\text{orb}(g, \tau)}, \tau)$ is isomorphism to a system of order d .

3.3. Tori. A d -dim torus is $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$. An integer-valued $d \times d$ matrix A can be seen as a homomorphism of $\mathbb{R}^d$ and $\mathbb{Z}^d$, and it induces a homomorphism on $\mathbb{T}^d$. A map $\tau : \mathbb{T}^d \rightarrow \mathbb{T}^d$ is an affine transformation if there are a matrix $A \in \mathbb{Z}^{d \times d}$ and a fixed point $b_0 \in \mathbb{T}^d$ such that $\tau(b) = Ab + b_0$.

Note that if the integer-valued $d \times d$ matrix A is a zero entropy homomorphism, then all the eigenvalues of A are unity roots. In fact, according to Theorem 2.1, if the integer-valued D is a zero entropy homomorphism, then $\prod_{j=1}^d \lambda_j$ is a non zero integer with all $|\lambda_j| \leq 1$, which means $|\lambda_j| = 1$. And then by the following Kronecker Lemma we have that all the λ_j are unity roots.

Theorem 3.3 (pp 108, Lemma (a)). [10] *If $\{\lambda_j\}_{j=1}^d$ satisfies $|\lambda_j| = 1$, all the j , and*

$$p(x) = \prod_{j=1}^d (x - \lambda_j)$$

are integral coefficient polynomial, then all the λ_j are unity roots.

Let

$$\rho(x, y) = \max_{1 \leq i \leq d} |x_i - y_i|, \quad x = (x_j)_j, y = (y_j)_j \in \mathbb{T}^d$$

be a metric of $\mathbb{T}^d$. Now we begin to consider the return time $N(a, \tau, \epsilon)$ of arbitrary $a = (a_j)_j \in \mathbb{T}^d$, where $N(a, \tau, \epsilon) = \{n \in \mathbb{Z} : \rho(\tau^n a, a) < \epsilon\}$.

Proposition 3.4. *Let $X = \mathbb{T}^d$, τ be a affine transformation of X with zero entropy, and $a = (a_j)_j \in X = \mathbb{T}^d$. Then for any ϵ , $N(a, \tau, \epsilon)$ include some d -degree generalized polynomial return time. i.e.*

$$N(a, \tau, \epsilon) \supset \bigcup_{r=0}^{b-1} \left(\bigcap_{j=1}^d \{tb + r, t \in N_{rj}\} \right),$$

where each N_{rj} is a return time of a polynomial function f_{rj} such that $N_{rj} = \{t, |\{f_{rj}(t)\} - a_j| < \epsilon\}$ for polynomials f_{rj} with degree less than $d + 1$.

Proof. Let $\tau : \mathbb{T}^d \rightarrow \mathbb{T}^d$, $(x_j)^T \rightarrow A(x_j)^T + (\alpha_j)^T$ be an affine transformation with zero entropy. Where $\alpha = (\alpha_j) \in \mathbb{T}^d$ is a fix point. Since τ has zero entropy, then A is a $d \times d$ integral matrix. Since Kronecker Lemma, all the eigenvalues of A is roots of unity. That means there are some B and P such that $A = P^{-1}BP$, where B is a upper triangular matrix, and all the entries in the diagonal of B are roots of unity.

First we have

$$\begin{aligned} \tau^n a &= A^n(a_j)^T + \sum_{i=0}^{n-1} A^i(\alpha_j)^T \\ &= P^{-1}B^n P(a_j)^T + P^{-1}\left(\sum_{i=0}^{n-1} B^i\right)P(\alpha_j)^T. \end{aligned}$$

Since B is a upper triangular matrix, and all the diagonal elements of B is roots of unity, there is b such that B^b is a upper triangular matrix with all the diagonal elements is 1. Denote $C = B^b$. Now for $n = tb + r$, $0 \leq r \leq b - 1$,

$$\tau^n a = P^{-1}C^t B^r P(a_j)^T + P^{-1}\left(\sum_{i=0}^{b-1} B^i\right)\left(\sum_{k=0}^{t-1} C^k\right)P(\alpha_j)^T + P^{-1}\left(\sum_{i=0}^{r-1} B^i\right)C^t P(\alpha_j)^T.$$

For the fix r , we consider if $\rho(a, \tau^{tb+r} a) < \epsilon$. Let $C = I + D$, here D is strictly upper triangular matrix, $D^d = 0$. Then $C^t = \sum_{i=0}^{d-1} \frac{t!}{i!(t-i)!} D^i$. That means each element of C^t is a polynomial of t , since r is fixed, each element of $P^{-1}C^t B^r P$, $P^{-1}(\sum_{i=0}^{b-1} B^i)(\sum_{k=0}^{t-1} C^k)P$, $P^{-1}(\sum_{j=0}^{r-1} B^j)C^t P$ is a polynomial of t ; that means each coordinate of $\tau^{tb+r} a$ is a polynomial of t with degree no more than a fix $d \in \mathbb{N}$:

$$\tau^{tb+r} a = (f_{rj}(t))_{1 \leq j \leq d}.$$

And all f_{rj} are no more than d degree. So for $N_{rj} = \{t, |\{f_{rj}(t)\} - a_j| < \epsilon\}$, here $\{f_{rj}(t)\} = f_{rj}(t) \pmod{1}$,

$$N(a, \tau, \epsilon) = \bigcup_{r=0}^{b-1} \left(\bigcap_{j=1}^m \{tb + r, t \in N_{rj}\} \right).$$

The proof is completed. $\square$

3.4. Nilpotent Lie groups. Now we begin to prove Theorem 3.1 for the nilpotent Lie group case.

Proposition 3.5. *Let A be a continuous surjective homomorphism of k -step nilpotent Lie group G , and let $G_i = [G_{i-1}, G]$ with $G_0 = G$. Then A induces naturally surjective homomorphisms A_i :*

$$A_i : G_i/G_{i+1} \rightarrow G_i/G_{i+1},$$

and A_i is invertible linear mapping.

We call $\{A_1, A_2, \dots, A_k\}$ the linear part of A .

Since each nilpotent Lie group can be imbedded into a upper triangular matrix group, in this paper, we only analysis the matrix nilpotent Lie group.¹ We could write a element of k -step matrix nilpotent Lie group G as:

$$\begin{pmatrix} 1 & x_{1,1} & x_{2,1} & \dots & x_{k-1,1} & x_{k,1} \\ 0 & 1 & x_{1,2} & \dots & \dots & x_{k-1,2} \\ 0 & 0 & 1 & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & 1 & x_{1,k} \\ 0 & \dots & \dots & \dots & 0 & 1 \end{pmatrix}$$

In the following part of this paper, we write a element of G as $(x_{i,j})_{1 \leq i \leq k, 1 \leq j \leq k+1-i}$, or just write it as $(x_{i,j})_{i,j}$ if there is no confusion. Group operation is the matrix operation. And we write the element of G_i/G_{i+1} as $(x_{i,1}, x_{i,2}, \dots, x_{i,k+1-i})$ or $\vec{x}_i$. Here we would define a new kind of degree of a polynomial. The polynomial is define on the matrix elements of of a element of nilpotent Lie group G .

Definition 3.6. For $(x_{i,j})_{1 \leq i \leq k, 1 \leq j \leq k+1-i}$, let

$$W = \left\{ \prod_{1 \leq i \leq k, 1 \leq j \leq k+1-i} x_{i,j}^{\alpha_{i,j}} : \alpha_{i,j} \in \mathbb{N} \right\}.$$

And define L -polynomial degree $l : W \rightarrow \mathbb{N}$ such that

$$l\left(\prod_{1 \leq i \leq k, 1 \leq j \leq k+1-i} x_{i,j}^{\alpha_{i,j}}\right) = \sum_{1 \leq i \leq k, 1 \leq j \leq k+1-i} 2^{i-1} \alpha_{i,j}$$

If f is a constant, $l(f) = 0$. For the polynomial f defined on the terms of $(x_{i,j})_{1 \leq i \leq k, 1 \leq j \leq k+1-i}$, $f = c_1 w_1 + \dots + c_t w_t$, $c_i \in \mathbb{R}$, $w_i \in W$, we define $l(f) = \max_{i=1,2,\dots,t} l(w_i)$. Also it is clear that $l(fg) \leq l(f) + l(g)$ and $l(f+g) \leq \max\{l(f), l(g)\}$.

¹We can see this conclusion from Ado theorem. Also we know that each G/Γ can be seen as a submanifold of $\hat{G}/\hat{\Gamma}$ such that $\hat{G}$ is connect Lie group. Then we use the linear property of a Lie algebra and transformation between Lie algebra and Lie group to see this conclusion.

Here we give the conclusion of matrix nilpotent Lie group G .

Proposition 3.7. *Let G be a k -step matrix nilpotent Lie group. A is a surjective continuous homomorphism of G . For any $g = (x_{i,j})_{1 \leq i \leq k, 1 \leq j \leq k+1-i}$, $A((x_{i,j})_{i,j}) = (\varphi_{i,j}(g))_{i,j}$, we have:*

(3.1)

$$\varphi_{i,j}(g) = A_{i,j}(x_{i,1}, x_{i,2} \dots x_{i,k+1-i}) + f_{i,j}(x_{1,1}, \dots x_{1,k}, x_{2,1} \dots x_{2,k-1} \dots x_{i-1,1} \dots x_{i-1,k+2-i})$$

where $A_{i,j}$ is linear functional, $f_{i,j}$ is a polynomial such that its L -polynomial degree $l(f_{i,j})$ is no more than 2^{i-1} .

Remark 3.8. We might use Lie algebra to prove a similar proposition. But to make the degree more exact, we use a complex proof.

Proof. It is clear that for each G_i , if A is surjective, $AG_i = G_i$. Then the first part of (3.1) is clear. (i.e. $A_{i,j}$ is linear.)

Now we would prove that $f_{i,j}$ is polynomial with $l(f) \leq 2^{i-1}$. Because A is homomorphism, $A(gg') = A(g)A(g')$. we have:

$$(\varphi_{i,j}(g))(\varphi_{i,j}(g')) = (\varphi_{i,j}(gg')).$$

That means

$$(3.2) \quad \varphi_{i,j}(gg') = \sum_{s=0}^i \varphi_{s,j}(g) \varphi_{i-s,s+j}(g').$$

where $\varphi_{0,j}(g) = 1$.

Let $\{1, \dots, k\}$ act on G such that $t(g)$ be the matrix in G and

$$t(g) = (x_{i,j,t})_{i,j} \quad x_{t,j,t} = x_{t,j}, \quad x_{i,j,t} = 0 \text{ if } i \neq t.$$

we have $t(g) \in G_t$, $g = 1(g)2(g) \dots k(g)$, and let $h_t(g) = 1(g) \dots t(g)$, $h_0(g) = I$. Since $AG_i = G_i$, $\varphi_{i,j}(t(g)) = 0$ if $t > i \geq 1$. Then (3.2) gives the result that for $i > 0$

$$\varphi_{i,j}(h_{t+1}(g)) = \sum_{s=0}^i \varphi_{s,j}(h_t(g)) \varphi_{i-s,s+j}((t+1)(g)) = \varphi_{i,j}(h_t(g)) \text{ if } t \geq i.$$

Which means $\varphi_{i,j}(g) = \varphi_{i,j}(h_i(g))$. And we have:

$$\begin{aligned} & \varphi_{i,j}(h_i(g)) \\ &= \sum_{s=0}^i \varphi_{s,j}(h_{i-1}(g)) \varphi_{i-s,s+j}(i(g)) \\ &= \varphi_{i,j}(h_{i-1}(g)) + \varphi_{i,j}(i(g)) \\ (3.3) \quad &= \sum_{s=0}^i \varphi_{s,j}(h_{i-2}(g)) \varphi_{i-s,s+j}((i-1)(g)) + \varphi_{i,j}(i(g)) \\ &= \varphi_{i,j}(h_{i-2}(g)) + \varphi_{i-1,j}(h_{i-2}(g)) \varphi_{1,j}((i-1)(g)) + \varphi_{i,j}((i-1)(g)) + \varphi_{i,j}(i(g)) \\ &= \dots \\ &= \sum_{s=1}^i \varphi_{i,j}(s(g)) + \sum_{s=1}^{i-1} \sum_{r=1}^s \varphi_{i-r,j}(h_{i-1-s}(g)) \varphi_{r,j-r}((i-s)(g)). \end{aligned}$$

For $i = 1$, $l(f_{1,j}) \leq 1$ since G/G_1 is a linear space. (3.1) is right for $i = 1$. Now we make the induction for $1, 2, \dots, i-1$. i.e. for $s \leq i-1$,

$$l(\varphi_{s,j}(g)) = l(\varphi_{s,j}(h_s(g))) \leq 2^{s-1}$$

By the induction, $l(\varphi_{i-r,j}(h_{i-1-s}(g)) \varphi_{r,j-r}((i-s)(g))) \leq 2^{i-r-1} * 2^{r-1} \leq 2^{i-1}$.

Claim: $\varphi_{i,j}(s(g))$ is a polynomial on $x_{i,j}$ with $l(\varphi_{i,j}(s(g))) \leq 2^{i-1}$, where $s \leq i$.

If the claim is right, according to and (3.3), $\varphi_{i,j}(h_{i-1}(g))$ satisfies $l(\varphi_{i,j}(h_{i-1}(g))) \leq 2^{i-1}$. Then $l(\varphi_{i,j}(g)) \leq 2^{i-1}$. Hence this proposition is right.

Now we prove the claim. For a fix s , we denote $(y_1, y_2, \dots, y_{x+1-s})_s$ being the element $\hat{g} = (\hat{x}_{i,j})_{i,j} \in G$ such that

$$(\hat{x}_{s,1}, \dots, \hat{x}_{s,k+1-s}) = (y_1, y_2, \dots, y_{k+1-s}) \text{ and } x_{i,j} = 0 \text{ for } i \neq s.$$

According to the operation of matrix, $(0, 1, 0, \dots, 0)_s * (1, 0, \dots, 0)_s = (1, 1, 0, \dots, 0)_s$, here each coordinate corresponding a matrix. What is more, we have:

$$(y_1, y_2, \dots, y_{k+1-s})_s = (0, \dots, 0, y_{k+1-s})_s * (0, \dots, 0, y_{k,s}, 0)_s \dots, (y_1, 0, \dots, 0)_s.$$

For $1 \leq s \leq k$, denote $\gamma_{r,s} = (0, \dots, 0, x_r, 0, \dots, 0)_s \in G$ corresponding to $g = (x_{i,j})_{i,j} \in G$. Then $s(g) = \gamma_{k+1-s,s} \dots \gamma_{1,s}$. By (3.2),

$$\varphi_{i,j}((x_1, x_2, \dots, x_{k+1-s})_s) = \sum_{r=0}^i \varphi_{r,j}(\gamma_{k+1-s}) \varphi_{i-r,j+r}((x_1, x_2, \dots, x_{k-s}, 0)_s).$$

According to the induction, if we want to prove $l(\varphi_{i,j}((x_1, x_2, \dots, x_{k+1-s})_s)) \leq 2^{i-1}$, we only need to prove

$$l(\varphi_{i,j}((x_1, x_2, \dots, x_{k-s}, 0)_s)) \leq 2^{i-1}$$

and

$$l(\varphi_{i,j}(\gamma_{k+1-s})) \leq 2^{i-1}.$$

We use (3.2) again and again for $s(g) = \gamma_{k+1-s,s} \dots \gamma_{1,s}$, then we just need to prove $l(\varphi_{i,j}(\gamma_r)) \leq 2^{i-1}$ for each r . Let $e_r = (0, \dots, 0, 1, 0, \dots, 0)_s$. Then e_r generated a one-dim sub Lie group of G . We know $e_r^n = ne^r$ for positive integer n , which means

$$A(ne_r) = A(e_r^n) = (A(e_r))^n.$$

$\gamma_r = e_r^{x_r}$, where e_r^x is defined by $q_n \in \mathbb{Q}, q_n \rightarrow x$ such that $e_r^x = \lim_{n \rightarrow \infty} e_r^{q_n}$, and $e_r^{\frac{1}{t}}$ is the matrix C satisfying $C^t = e_r$. For $e_r \in G$, the matrix C is unique.

Then since A is continuous, $(\varphi_{i,j}(\gamma_r))_{i,j} = A(e_r^{x_r}) = (Ae_r)^{x_r}$. Then for $(Ae_r)^{x_r}$, since $B = A(e_r) = (b_{i,j})_{i,j} \in G$, then each matrix element of B^{x_r} is a function of x_r . What is more, since the diagonal element of B is 1, $\varphi_{i,j}(\gamma_r)$ is a polynomial of x_r . Also $b_{t,j} = 0$ for $1 \leq t < s$. Since $A(G_s) \subset G_s$, we know that the degree x_r in the polynomial $\varphi_{i,j}(\gamma_r)$ of x_r is no more than 2^{i-s} , which means that $l(\varphi_{i,j}(\gamma_r)) \leq 2^{i-1}$.

So the claim is right. The whole proof is completed. $\square$

Now for the nilpotent manifold G/Γ and its zero entropy affine transformation, we prove Theorem 3.1.

For an affine transformation of G/Γ , $\tau : g \rightarrow \alpha A(g)$ with $\alpha \in G$ and A being an automorphism of G/Γ , which induced by a homomorphism A of G . A also keep Γ . τ has zero entropy if and only if A has zero entropy. In this case, each A_i in Proposition 3.5 is zero entropy (each A_i also induce a automorphism on G_i/Γ_i).

And for each i , we have $A_i = Q_i^{-1} B_i Q_i$, where B_i is a upper triangular matrix and all the diagonal elements of B_i are roots of unity. Also there is a matrix P such that $B = P^{-1} A P$ with B_i is the linear part of B . Let b be the integer such that all the diagonal elements of B_i^b is 1 for each i . Let $C = A^b$. Then for n , we have $n = tb + r$ and $A^n = C^t A^r$ where $0 \leq r \leq b - 1$.

3.5. Proof of Theorem 3.1. We need a metric of G/Γ which can generate the topology. One simple metric ρ is the maximum distance such that for $g\Gamma, g'\Gamma \in G/\Gamma$ with $g = (x_{i,j})_{i,j}$ and $g' = (x'_{i,j})_{i,j}$ $x_{i,j}, x'_{i,j} \in [0, 1)$, $\rho(g\Gamma, g'\Gamma) = \max_{i,j} |x_{i,j} - x'_{i,j}|$. For each $g\Gamma \in G/\Gamma$, we only need to consider the open neighborhood $U_\epsilon = \{h + \Gamma, \rho(h\Gamma, g\Gamma) < \epsilon\}$.

We consider the affine transformation $\tau(g\Gamma) = g_0 A(g\Gamma)$ on G/Γ , where A is a zero entropy homomorphism on G (it induces a map on G/Γ) and fix element $g_0 \in G$. Also A satisfies $A\Gamma \subset \Gamma$. Here we denote $g_0 = (y_{i,j})_{i,j}$.

Let b be the integer such that all A_i (the linear part of A) satisfies that the eigenvalues of A_i^b are 1 (as the discussion in torus case).

Claim: Each matrix element of $A^{nb+r}(g) = (\phi_{i,j}(n, g))_{i,j}$ is a finite degree polynomial of n . The degree of $\phi_{i,j}(n, g)$ is less than some M , which is independent of g .

The claim is right since Proposition 3.7 and $A^{nb+r}(g) = (A^b)^n(g) A^r(g)$. We know that $B = A^b$ is still a group automorphism. If we write $B(g) = (\varphi_{i,j}(g))_{i,j}$, then

$$\varphi_{i,j}(g) = B_{i,j}(x_{i,1}, x_{i,2} \dots x_{i,k+1-i}) + h_{i,j}(x_{1,1}, \dots, x_{1,k}, x_{2,1} \dots x_{2,k-1} \dots x_{i-1,1} \dots x_{i-1,k+2-i})$$

Where $A_i^b = B_i = (B_{i,1} \dots, B_{i,k+1-i})$, which means that B_i is the matrix with all eigenvalues being 1. We write $h_{i,j}(x_{1,1}, \dots, x_{1,k}, x_{2,1} \dots x_{2,k-1} \dots x_{i-1,1} \dots x_{i-1,k+2-i}) = H_{i,j}(g)$, $(x_{i,1}, x_{i,2} \dots x_{i,k+1-i}) = \vec{x}_i$, $\vec{H}_i(g) = (H_{i,1}(g), \dots, H_{i,k+1-i}(g))$. Then

$$(\phi_{i,1}(n), \dots, \phi_{i,k+1-i}(n)) = B_i^n(\vec{x}_i) + \sum_{j=0}^{n-1} B_i^{n-1-j}(\vec{H}_i(B^j g))$$

We make an induction for i to show that $\phi_{i,j}(n)$ are polynomials of n .

$i = 1$ is clear since the tori case. Now we assume that for $1 \leq t \leq i-1$, $\phi_{t,j}(n)$ are polynomials of n . Then $H_{i,j}(B^m g)$ are polynomials of m since $h_{i,j}(\ast)$ are polynomials and $H_{i,j}(B^m g)$ is a polynomial of $\phi_{t,j}(m)$ with $1 \leq t \leq i-1$.

Also if $\vec{z} = (z_1, \dots, z_{k+1-i})$ is fixed, each coordinate of $B_i^m(\vec{z})$ are polynomials of m since the discussion of tori. what is more, for fix m , each coordinate of $B_i^m(z_1, \dots, z_{k+1-i})$ is a linear combination of $z_1, \dots, z_{k+1-i}$. That means, each coordinate of $B_i^{n-1-j}(\vec{H}_i(B^j g))$ are polynomials of n and j . Then each coordinate of $B_i^{-j}(\vec{H}_i(B^j g))$ are polynomials of j , we write it as $B_i^{-j}(\vec{H}_i(B^j g)) = (v_1(j), \dots, v_{k+1-i}(j))$. Where $v_s(j)$ are polynomials of j .

Then

$$\sum_{j=0}^{n-1} B_i^{n-1-j}(\vec{H}_i(B^j g)) = B_i^{n-1} \left(\sum_{j=0}^{n-1} v_1(j), \dots, \sum_{j=0}^{n-1} v_{k+1-i}(j) \right)$$

Since for arbitrary fix $m \in \mathbb{N}$, $\sum_{j=0}^{n-1} j^m$ are polynomials of n . Then $\sum_{j=0}^{n-1} v_s(j)$ are polynomials of n for $1 \leq s \leq k+1-i$. Then we have $\phi_{i,j}(n)$ are polynomials of n for each j . That is the claim.

Now we consider the zero entropy affine τ .

$$\tau^{nb+r}(g\Gamma) = A(g_0) \dots A^{nb+r-1}(g_0) A^{nb+r}(g_0) A^{nb+r}(g)\Gamma.$$

We consider $A(g_0) \dots A^{nb+r-1}(g_0) A^{nb+r}(g_0) A^{nb+r}(g)$. Notice that $A^m(g_0)$ is a upper triangle matrix with the diagonal element being 1. The claim tells us that

$A^{(n-1)b+r+1}(g_0) \dots A^{nb+r}(g_0)$ is a upper triangle matrix with diagonal element being 1 that each matrix element of it is a finite degree polynomial of n . Then according to matrix operation, we know that each matrix element of $A(g_0) \dots A^{nb+r-1}(g_0)A^{nb+r}(g_0)$ is a finite degree polynomial of n .

Now we denote $A(g_0) \dots A^{nb+r-1}(g_0)A^{nb+r}(g_0)A^{nb+r}(g) = (p_{i,j,r}(n))_{i,j}$, where each $p_{i,j,r}(n)$ is a polynomial with degree less than some M_0 independent with g . We want to find $g_{n,r}\Gamma = (p_{i,j,r}(n))_{i,j}\Gamma$ such that $g_n = (q_{i,j,r}(n))_{i,j}$ with $q_{i,j,r}(n) \in [0, 1]$.

For $i = 1$, $q_{1,j,r}(n) = p_{1,j,r}(n) \pmod{1}$. We want to find $\gamma_{n,r} = (y_{i,j,r}(n))_{i,j} \in \Gamma \subset M_{k+1}(\mathbb{Z})$ such that $g_n\gamma_{n,r} = (p_{i,j,r}(n))_{i,j}$, that means

$$p_{i,j,r}(n) = \sum_{s=0}^i q_{s,j,r}(n)y_{i-s,s+j,r}(n).$$

Here $q_{0,j',r} = 1, y_{0,j',r} = 1$ for all j' . Then $y_{1,j'} = [p_{1,j'}]$ are finite degree generalized polynomial for all j' .

Inductively, we assume $q_{1,j',r}, \dots, q_{i-1,j',r}, \gamma_{1,j',r}, \dots, \gamma_{i-1,j',r}$ are generalized polynomial with finite degree for all j' . Then

$$q_{i,j,r} + y_{i,j,r} = p_{i,j,r} - \sum_{s=1}^{i-1} q_{s,j,r}y_{i-s,s+j,r}$$

Since $\gamma_{n,r} \in \Gamma$,

$$y_{i,j,r} = [p_{i,j,r} - \sum_{s=1}^{i-1} q_{s,j,r}y_{i-s,s+j,r}]$$

while

$$q_{i,j,r} = \{p_{i,j,r} - \sum_{s=1}^{i-1} q_{s,j,r}y_{i-s,s+j,r}\}$$

are generalized polynomials with finite degree independent with g .

That is what we want:

$$N(g\Gamma, U_\epsilon) = \bigcup_{r=1}^b \left(\bigcap_{i,j} \{nb+r : |q_{i,j,r}(n) - q_{i,j,r}(0)| < \epsilon\} \right)$$

Then the theorem is right for matrix nilmanifold.

3.6. Proof of Theorem 3.2. The integer b is given earlier. For each $1 \leq r \leq b$, $(\overline{orb}(\tau^r g, \tau^b), \tau^b)$ is a dynamical system. Also we have $\overline{orb}(g, \tau) = \bigcup_{1 \leq r \leq b} \overline{orb}(\tau^r g, \tau^b)$. The reason is, for each $x \in \overline{orb}(g, \tau)$, if $x_j \rightarrow_{j \in J} x$ with $x_j \in \overline{orb}(g, \tau)$, then there is r_x such that $\{x_j\}_{j \in J} \cap \overline{orb}(\tau^{r_x} g, \tau^b)$ are infinite set. Let $X_j = \overline{orb}(\tau^{r_x} g, \tau^b)$.

According to the proof of Theorem 3.1 and Theorem 2.6, (X_r, τ^b) are systems which are AA system. According to Theorem 2.4, we know that the system is a almost one to one extension of its maximal nilfactor (Y_r, S_r) with factor map π_r . Since (X_r, τ^b) is minimal and each point $x \in X_r$ is AA point, it is one to one extension. i.e. $(Y_r, S_r) = (X_r, \tau^b)$.

Now we construct a dynamical system (Y, S) . We consider each Y_r as different dynamical systems with national isomorphic map $I_r : Y_r \rightarrow Y_{r+1}$. (The map is induced by $\tau : X_r \rightarrow X_{r+1}$). Also $I_d : Y_d \rightarrow Y_1$ is induced by $\tau : X_d \rightarrow X_1$. Let

$Y = \bigsqcup_{r=1}^b Y_r$, $S|_{Y_r} = I_r$. Since each (Y_r, S^b) is a system of order d , by the property of nilsystem, (Y, S) is a system of order d . Since $(\overline{\text{orb}(g, \tau)}, \tau)$ is a factor of (Y, S) , it is a system of order d .

3.7. Compact abelian groups. We also prove a similar but weaker conclusion for abelian compact group. That is:

Theorem 3.9. *Let X be a abelian compact group, and $\tau : X \rightarrow X, x \mapsto \alpha x x_0$ be an affine transformation with zero entropy. Where α is zero entropy automorphism. Then for each $x \in X$, $(\overline{\text{orbit}(x, \tau)}, \tau)$ is isomorphic to an inverse limit of nilsystems.*

Now let us consider a compact abelian group X with zero entropy affine transformation.

The affine transformation of X is $\tau : X \rightarrow X$, $\tau(g) = A(g)g_0$. Here A is a continuous automorphism, $g_0 \in X$. However, since we don't need the A -invariant metric and each system has an invertible fact, we can also use Property 3.10 for A . And we need a property.

Property 3.10. [18] *Let α be an automorphism of a compact group X with zero entropy. Then there exists a decreasing sequence of closed, normal, α -invariant subgroups*

$$X = Y_0 \supset Y_1 \supset \dots \supset Y_n \supset \dots$$

such that $\bigcap_{i \geq 1} Y_i = \{1_X\}$, and Y_{i-1}/Y_i is a compact Lie group with an α^{Y_{i-1}/Y_i} -invariant metric for every $i \geq 1$.

In particular, there exists no infinite, closed, α -invariant subgroup $Y \subset X$ such that α^Y is ergodic.

Proof of Theorem 3.9

According to Property 3.10, for a compact abelian group X with zero entropy automorphism α , there exists a decreasing sequence of closed, normal, α -invariant subgroups

$$X = Y_0 \supset Y_1 \supset \dots \supset Y_n \supset \dots$$

such that $\bigcap_{i \geq 1} Y_i = \{1_X\}$, and Y_{i-1}/Y_i is a compact Lie group with an α^{Y_{i-1}/Y_i} -invariant metric for every $i \geq 1$.

For the zero entropy affine transformation τ of G , $\tau(x) = \alpha x x_0$. We notice that for the compact abelian Lie group $(X/Y_i, \alpha_i)$, where $\alpha_i : xY_i \rightarrow \alpha(x)x_0Y_i$ is a zero entropy affine transformation for X/Y_i , also the inverse limit of $(X/Y_i, \alpha_i)$ is X .

Also for each $x \in X$, $(\overline{\text{orb}(x, \tau)}, \tau)$ is the inverse limit of $(\overline{\text{orb}(xY_i, \tau_i)}, \tau_i)$. According to Theorem 3.2, we know that each $(\overline{\text{orb}(xY_i, \tau_i)}, \tau_i)$ is a system of order d_i . Then $(\overline{\text{orb}(x, \tau)}, \tau)$ is a ∞ -order system.

REFERENCES

- [1] V. Bergelson, B. Host and B. Kra, *Multiple recurrence and nilsequences. With an appendix by Imre Ruzsa*, Invent. Math. 160 (2005), no. 2, 261–303.
- [2] V. Bergelson, J. Rosenblatt, *Mixing actions of groups*, Illinois J. Math. 32 (1988), no. 1, 65C–80.
- [3] N.P. Brown, *Topological entropy in exact C^* -algebras*, Math. Ann. 314 (1999), 347–367

- [4] L. J. Corwin and F. P. Greenleaf, *Representations of nilpotent Lie groups and their applications*. Part I. Basic theory and examples, Cambridge Studies in Advanced Mathematic 18, Cambridge University Press, Cambridge, 1990. vii+269 pp.
- [5] P. Dong, S. Donoso, A. Maass, S. Shao and X.D. Ye, *Infinite-step nilsystems, independence and complexity*, Ergod. Th. & Dynam. Sys. (2013), 33, 118-143
- [6] R. Ellis and H.B. Keynes, *Bohr compactification and a result of Følner*, Israel J. Math., 12(1972), 314-30
- [7] H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, M. B. Porter Lectures. Princeton University Press, Princeton, N.J., 1981.
- [8] B. Green, T. Tao, *The Möbius function is strongly orthogonal to nilsequences*, Ann. Math. (2), 175 (2012), 541–566.
- [9] B. Hall, *Lie groups, Lie algebras, and representations. An elementary introduction*, Graduate Texts in Mathematics, 222. Springer-Verlag, New York, 2003. xiv+351 pp.
- [10] E. Hecke, *Lectures on the Theory of Algebraic Numbers*. Graduate Texts in Mathematics, 77, Springer-Verlag, New York.
- [11] B. Host, B. Kra, *Analysis of two step nilsequences*, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 5, 1407C-1453.
- [12] B. Host, B. Kra, *Nil-Bohr sets of integers*, Ergodic Theory Dynam. Systems, **31** (2011), no. 1, 113–142.
- [13] B. Host, B. Kra and A. Maass, *Nilsystèmes d'ordre deux et parallélépipèdes*, Bull. Soc. Math. France, 135 (2007) 367-405
- [14] B. Host, B. Kra and A. Maass, *Nilsequences and a structure theory for topological dynamical systems*, Adv. Math., 224 (2010) 103–129.
- [15] B. Host, A. Maass, *Nilsystèmes d'ordre deux et parallélépipèdes*, Bull. Soc. Math. France, 135 (2007) 367–405.
- [16] W. Huang, S. Shao and X.D. Ye, *Nil Bohr-sets and almost automorphy of higher order*, to appear in Memoirs of the AMS
- [17] J.Y. Liu, P. Sarnak, *The Möbius function and distal flows*, Duke Math. J. 164 (2015), no. 7, 1353–1399.
- [18] K. Schmidt, *Dynamical systems of algebraic origin*, Progress in Mathematics, 128. Birkhäuser Verlag, Basel, 1995. xviii+310 pp.
- [19] Y. Sinai, *On the concept of entropy for a dynamical system*, Dokl. Akad. Nauk SSSR 124 (1959), 768–771.

WU WEN-TSUN KEY LABORATORY OF MATHEMATICS, USTC, CHINESE ACADEMY OF SCIENCES AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, ANHUI, 230026, P.R. CHINA, AND DEPARTMENT OF MATHEMATICS, SUNY AT BUFFALO, BUFFALO, NY 14260-2900, U.S.A.

E-mail address: lianzx@mail.ustc.edu.cn, zhengxin@buffalo.edu